

White Paper: The One-Size-Fits-All Age Adjusted Maximum Heart Rate Equation Fits No One

Sally Edwards, MA, MBA, Author, Exercise Scientist, CEO Heart Zones, Inc.©

Position Statement

The 220-minus-age equation regularly used to estimate an individual's maximum heart rate (MHR) is invalid¹ for this purpose. The equation, first published in 1970 from data with large variance, disregards the range of individual differences. MHR in healthy adult individuals can be as low as 150 beats-perminute (BPM) and as high as 230 BPM. Because of this large variance, group data averaged to a mean cannot be used to predict the MHR of individuals with reasonable accuracy. Such use is inappropriate, inaccurate and can be dangerous if used to derive exercise prescriptions. There is no equation that accurately estimates an individual's MHR or any other human biomarker, including blood pressure or cholesterol levels. Rather, exercise experts that I work with recommend individual assessments using a sub-max field test to estimate individual MHRs. We further argue that models based on population averages usually lead to inaccurate estimates and therefore the MHR equals 220-minus-age equation should not be used.

Discussion:

By definition, MHR is the highest number of times per minute the heart can contract. It's the heart rate at the point of exhaustion in an all-out short effort. An individual maybe be able to continue briefly to increase effort or speed when the actual MHR number is reached, but the heart simply won't beat any faster. Most scientists think individual MHR is a genetically determined number and not based

¹ GELLISH, RONALD L.; GOSLIN, BRIAN R.; OLSON, RONALD E.; McDONALD, AUDRY; RUSSI, GARY D.; MOUDGIL, VIRINDER K. Longitudinal Modeling of the Relationship between Age and Maximal Heart Rate. Medicine & Science in Sports & Exercise. 39(5):822-829, May 2007. "Although the HRmax-prediction model of 220 - age has become well established in the medical literature and is used widely in clinical and fitness settings, its validity is uncertain." <u>http://www.ncbi.nlm.nih.gov/pubmed/1746858</u>

on body size.² Rather, MHR is a self-protecting mechanism. If the heart beats too fast, it doesn't have enough time between beats to fill the chambers adequately in order to fully contract and effectively pump the volume of blood demanded by the effort.

The 220 BPM minus an individual's age equation for calculating the MHR almost accidentally became the standard in cardiology and in fitness programs, used in medical, sports and research. Indeed, an entire industry has grown around the use of this equation, especially the wearable and exercise equipment-enabled heart rate monitors sold to individuals and by cardio equipment manufacturers. Though inappropriate, as it is an average and not an individual number, the MHR 220-minus-age equation continues to be popular today because it is easy and seems commonsensical. Many exercise gurus, equipment handbooks, and coaches recite the equation. When learning to use most cardio-exercise equipment like a treadmill or indoor cycle, often there is a series of questions and one of this is "How old are you?". No matter how much science discredits the MHR 220-minus-age equation, no matter how many research studies invalidate it, too many professionals and exercise equipment manufacturers continue to believe that one size fits all – that everyone of the same age has the same MHR. This is false. Every child, every adolescent, and every adult has her or his own specific MHR, which is unique. That MHR number is genetically not mathematically determined.

The MHR 220-minus-age equation can lead to potentially dangerous outcomes. Carl Foster, Ph.D. and the past president of the American College of Sports Medicine, knows this from personal experience. When Dr. Foster, now professor of Exercise Science at the University of Wisconsin-La Crosse, was 20 and running collegiate cross country, his coach demanded that he run near his MHR. The cross country coach used the MHR 220-minus-age equation - Carl's age, 20, which predicted an MHR of 200 BPM as the goal he should achieve. But when Dr. Foster tested himself – after all, he was an exercise science major – in an all out sprint, the highest number that he could ever achieve was160 BPM. He was humiliated, ridiculed by the running coach in front of his teammates. But in fact, at an all out sprint, Dr. Foster's genetically determined MHR was within a few heartbeats of 160 BPM. The MHR

² Batterham, A. M.; George, K. P.; Mullineaux, D. R. Relationship Between Heart Size And Body Dimensions: An Allometric Scaling Approach: 886. Medicine & Science in Sports & Exercise: May 1995 - Volume 27 - Issue 5 - ppg S158

220-minus-age equation had a 40 BPM error for Dr. Foster, an egregious error. Training at too high a heart rate for too long a period, resulting in overtraining, can lead to hormonal, nutritional, mental, emotional, muscular, neurological and other imbalances.

Physician and consultant to U.S. soccer team, Dr. Donald Kirkendall, Duke University Medical Center Sports Medicine Section, is another of the professionals who challenge the seemingly etched-instone belief in the 220 minus age equation. In one of his first experiences analyzing the data from a heart rate monitor was testing a member of the U. S. rowing team, he told The New York Times. Dr. Kirkendall asked the athlete for an all-out effort for six minutes. One rower, in his mid-20s, had an estimated maximum heart rate of 195 BPM (220 minutes his age of 25 years or 195 BPM. "His pulse rate hit 200 at 90 seconds into the test," Dr. Kirkendall said. "And he held it there for the rest of the test", a much longer time than would be possible even if his MHR was actually 200, which it wasn't – it was substantially higher. A local cardiologist, looking on in astonishment, told Dr. Kirkendall, "You know, there's not a textbook in the world that says a person could have done that." The purpose of this White Paper is to support rewriting the textbooks on the 220 minus age equation.

So, why is it important that we know an individual's MHR? Who created the MHR 220-minusage equation and made it a belief so strong that even scientific rigor has not yet corrected it? Why if it is known that the prediction model is inaccurate hasn't the scientific community corrected it? And, given the enormous range in MHRs between individuals of the same age and fitness, why would anyone use the 220 minus age equation today?

Why Do We Need to know Individual MHR?

The 220 minus age MHR prediction is based on group averages of individuals of the same age. It says that all 10-year olds have the same MHR of 220 minus 10, or an MHR of 210 BPM, and all 50year olds have the same MHR of 170 BPM or 220 - 50 years. According to this and most MHR formulas³, as you get older, your MHR declines one BPM per year. This simply is not true.


³ Robergs, Robert A. and Landwehr, Roberto. Prediction of Maximal Heart Rate. Journal of Exercise Physiology. Volume 5 Number 2 May 2002. http://bit.ly/1ThD0V

In medical, research and exercise applications, an accurate measure of MHR is important as it is used for exercise testing and prescriptions for some of the following reasons:

- Determining the intensity of the exercise during a workout is often expressed as a percentage of MHR. For example, the average target heart rate for the workout might be only 72% of maximum effort.
- It is used to assess the fitness level. For example, an individual is becoming fitter because he or she is achieving a higher percentage of MHR for a longer period of exercise time.
- It becomes a criterion for achieving maximal exertion: For example, the individual's MHR is 180 BPM and in a high intensity interval training session (HIIT) the individual reached 170 BPM three times.
- For prescribing individual training zones for safe exercise activities in both the healthy and unhealthy population.

In exercise and fitness applications, MHR is used in both training and racing. One of the most important reasons to assess one's MHR accurately and individually is to use that number as an anchor to set training zones. The chart in Figure 1, while based on an individual's 220 minus age equation estimated MHR, can be used once the MHR is determined by ignoring the age axis. Simply use the MHR for the individual. Using heart rate training zones allows the fitness enthusiast a way to easily determine relative, working heart rate zones or workout zones using a wearable heart rate monitor. By using a zone training method, an individual can work out simply using percentages of the MHR number.

Over the last quarter century, dozens of scientists have tried to demonstrate that there is a more accurate measure of




Figures 1. Using MHR To Set Training Zones Based On MHR Equation 220 Minus Age. (©FITNUSPOSTERS)

MHR by adding more than just age as a variable in the equation. There now are MHR equations based

on ethnicity, gender, fitness levels, weight and other factors.⁴ And yet, there simply is no way to mathematically determine something that needs to be measured individually.⁵ Among practitioners who agree is exercise medicine researcher Mark Sarzynski's, PhD, Assistant Professor of Exercise Science, University of South Carolina. "Our findings show that based on the standard error of estimates, the prevailing age-based, estimated HR max equations do not precisely predict an individual's measured MHR."⁶

How is Maximum Heart Rate Determined?

There are two common ways to measure the individual MHR. The first way is to estimate MHR in either an all out effort or with sub-max testing protocols. The second but erroneous way to assess MHR for the individual is to use one of the many formulas from population averages. Once the MHR is



Figures 2. Heart Zones Calculated On 10% Of Measured Maximum Heart Rate (©Heart Zones, Inc)

⁴ J Am Coll Cardiol. 2001;37(1):153-156. doi:10.1016/S0735-1097(00)01054-8

⁵ M.A. SARZYNSKI,1, T. RANKINEN,1 C.P. EARNEST,2 A.S. LEON,3 D.C. RAO,4 J.S. SKINNER,5 and C. BOUCHARD1. Measured Maximal Heart Rates Compared to Commonly Used Age-Based Prediction Equations in the Heritage Family StudyAm J Hum Biol. 2013 Sep-Oct; 25(5): 695–701.

⁶ http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3935487/

determined, that number in BPM can then be used as an anchor point or value to calculate individual heart rate training zones, which are typically marked in 10% intervals of the MHR number.

The Maximum Heart Rate chart herein ⁷ shows five training zones each based on 10 per cent intervals of one's individual MHR.⁸ First published in 1993, <u>The Heart Rate Monitor Book ⁹</u>, this chart removes all factors used in the earlier equations such as age or gender. By anchoring training zones on one's actual individual MHR and setting goals based on percentages of a valid MHR number, a person or trainer can more accurately assess the appropriate amount of time to allot to each zone, arguably providing improved health and fitness benefits.

Where did the MHR equals 220-minus-age equation originate?

In 1927 Harvard University created the prestigious Harvard Fatigue Lab¹⁰ dedicated to promoting scientific and collaborative research in

exercise physiology. Then doctoral candidate Sid Robinson, who later became a professor at University of Indiana, published a paper that showed the MHR of different age groups. This is called cross-sectional versus longitudinal research. Robinson compared a group of 20year olds with a group of 30-year olds and so forth. The fitness levels were not measured nor were individuals screened for medication usage or health issues. Robinson's results appeared like a splattergram of dots because the individual




Figure 3. Splattergram of red dots are individual MHRs. The blue line is the regression line for the one-size-fits all equation. (Roberg)

⁷Heart Zones Inc copyright 1993

⁸ Carl Foster, The Bloodless Lactate Profile. Medicine and Science in Sports and Exercise, pp 927-933, 1995.

⁹ S. Edwards. The Heart Rate Monitor Book. Polar Electro Oy, 1993.

¹⁰<u>http://www.humankinetics.com/excerpts/excerpts/harvard-fatigue-laboratory-influential-in-promoting-exercise-physiology-research</u>

differences (i.e., variance) in MHR are so immense.¹¹ He and researchers who followed took the average of each age group regardless of fitness level, plotted it and drew a regression line that showed that MHR drops by 1 BPM per year. Today, longitudinal research does show that measured MHR may drop by age, especially in those who become unfit, but the rate of decline of MHR is individualistic, and that in large part as fitness declines the ability to reach MHR declines. That is the individual being tested may not have the strength or the stamina to reach their true MHR in a testing situation. Indeed, this decline in individual MHR may be the result of their individual inability to reach MHR rather than the MHR actually declining. There is currently no known research to assess this decline in MHR-testing capacity in the older population.

But, where did the one-size-fits all equation, the MHR 220-minus-age, originate? In 1970, three exercise medicine specialists in the Public Health Service, Fox, Naughton, and Haskell first derived and published it as a solution to a problem.¹² At the time, Dr. Samuel Fox was leading a program on heart health and was mentoring Dr. William Haskell. Together, with researcher J.P. Naughton, they were in search for a way to prescribe safe exercise intensities for cardiac patients.

But, according to Robert A. Robergs and Roberto Landwehr in the *Journal of Exercise Physiology*, "Surprisingly, there is no published record of research for this equation." It was not the Fox team's intention to create something that would become the foundation of aerobic exercise intensity levels, they simply sought a quick way to establish levels of effort in order to prescribe safe exercise. They were working with sick patients, not kids, fitness-seekers, or athletes and they needed to provide these patients with safe exercise prescriptions. Today there are a number of different equations¹³ ¹⁴ and the argument continues concerning which is the best equation with the least amount of error. In fact, all such equations to mathematically calculate MHR are equally useless. "I've kind of laughed about it over the years," Dr. William Haskell told The New York Times in 2001. The equation, he said,

¹¹ Robergs, Robert A. and Landwehr, Roberto. Prediction of Maximal Heart Rate. Journal of Exercise Physiology. Volume 5 Number 2 May 2002.

¹² Fox SM, III, Naughton JP, Haskell WL. Physical activity and the prevention of coronary heart disease. Ann Clin Res. 1971;3:404–432. [

¹³ Robergs RA, Landwehr R. The surprising history of the "HRmax=220 minus age" equation. J Exerc Physiol online. 2002;5:1–10.

¹⁴ Nes BM1, Janszky I, Wisløff U, Støylen A, Karlsen T. Age-predicted maximal heart rate in healthy subjects: The HUNT fitness study. Scand J Med Sci Sports. 2013 Dec;23(6):697-704. 2012 Feb 29.

"was never supposed to be an absolute guide to rule people's training." (For a more complete story about how the MHR equal 220 minus age equation became "perceived wisdom" read the entire New York Times article.¹⁵)

Why Do Some Individuals have a Higher MHR than Others of the Same Age?

Given identical age, fitness level, gender and all other factors, how can non-smokers who are disease and medication-free individuals have as high as an 80 BPM difference in their tested MHRs. A perhaps whimsical answer lies in the differences between the heart rate of a hummingbird versus that of a blue whale.

First, let's take the hummingbird's heart beat frequency - more than 20 times each second - or as high as 1,260 beats per minute. A hummingbird's heart is tiny, smaller than the size of an aspirin. Next, compare a hummingbird's heart to the world's largest heart - that of the seven-ton blue whale. A blue whale's heart can be as big as a room. The cardiac frequency of hummers and whales varies from individual to individual just like humans, but in general, larger animals have a slower heat rate than smaller animals. The average heart rate of large whales is from about 10 BPM to a MHR of 30 BPM.

But, a more technical answer to the question why there is such an enormous range in MHR between two individuals of the same weight, size, and other factors has to do with heart mass. The bigger the dimensions, the size and weight of your heart, the lower your MHR – you are a blue whale. And conversely, the smaller your heart mass is as compared to your body size, the higher your MHR – you are a hummingbird.

What Can Be Done to change the Practice of Using the MHR Equals 220-Minus-Age Equation?

There is a better way to estimate MHR than erroneously using a one-size fits all equation, which recent research has shown is especially inappropriate for women¹⁶ and children¹⁷. Our recommendation is to estimate an individuals MHR using a safe and time-efficient sub-max field test.

¹⁵ <u>http://www.nytimes.com/2001/04/24/health/maximum-heart-rate-theory-is-challenged.html?pagewanted=all</u>

¹⁶ Circulation. 2010 Jul 13;122(2):130-7. doi: 10.1161/CIRCULATIONAHA.110.939249. 2010 Jun 28. Heart rate response to exercise stress testing in asymptomatic women: the st. James women take heart project. Gulati M1, Shaw LJ, Thisted RA, Black HR, Bairey Merz CN, Arnsdorf MF.

¹⁷Dev Med Child Neurol. 2011 Sep;53(9):861-4. doi: 10.1111/j.1469-8749.2011.03989.x. Epub 2011 May 13.Maximum heart rate for children: The 220-age equation does not predict maximum heart rate in children and adolescents. http://www.ncbi.nlm.nih.gov/pubmed/21569015



Figure 4. The Sub Max Field Test To Estimate The Individual Maximum Heart. (@Heart Zones, Inc)

A field test or field experiment is different from a lab test. The field test is conducted in the same use environment or conditions of normal exercise instead of in a controlled environment like a laboratory. There are several different sub-max tests and when done in combination, these field tests are almost as accurate as an all-out MHR test.¹⁸ See the example of one of the "Easy-Moderate-Hard Sub-Max" field tests presented here in Figure 3.

Let's use the Easy-Moderate-Hard Sub Max Test as an example. After an adequate warm-up, the participant begins the first of three stages of an activity of their choice at an effort level that matches a feeling of effort best described as "Easy". After two minutes the participant records the number of beats-per-minute (BPM) that matches the feeling of Easy. Using the RPE, Rating of Perceived Exertion scale of 0 being no effort and 10 representing a feeling of all-out effort, the Easy level of intensity is a rating of 2. The second stage of the test is to immediately increase the intensity of the same activity to a moderate level that would be best represented on the RPE between a 4 to a 6 rating effort level. Again, after two minutes of moderate effort, the participant record their BPM for Moderate. The final 2-minute stage is at a hard to very hard level of intensity equal to a 6-7 on the scale of perceived exertion. Taking the BPM from all three stages of the Sub-Max Field Test the resulting estimate of MHR for this participant would be as 185 BPM. See Table 1.

Table 1. Calculating MHR from the Easy-Moderate-Hard Sub-Max Field Test							
Current Effort Level:	Easy	Moderate	Hard				
	120 BPM	145 BMP	170 BPM				
Add the Math Factor:	+60	+40	+20				
Estimated MHR from Field Test:	180 BPM	185 BPM	190 BPM	MHR: 185 BPM			

Convert from MHR to Threshold Heart Rate

There is another way to anchor individual training zones which is not based on maximum heart rate. This method is called the threshold heart rate method. Many exercise scientists and researchers today recommend this method of assessing for threshold heart rates because as your fitness level changes so does your threshold heart rate numbers. Threshold heart rate numbers are dynamic while the MHR number does not change with fitness.

¹⁸ Ten different Sub-Max tests are available in the Heart Rate Monitor Guidebook by Edwards. Heart Zones publishing 2010 available at <u>www.HeartZones.com</u>

Threshold biomarkers are measured when there is a shift in the energy requirements, the carbohydrates and fats, that are burned during an exercise session. The most accurate way to assess the threshold biomarkers is in an exercise laboratory while field tests for the threshold heart rate have been scientifically validated.¹⁹ There are two different threshold exercise intensities called the first threshold (T1) or the low threshold. The second threshold (T2) or the high threshold is commonly called the ventilatory or lactate threshold and formerly known as the anaerobic threshold. By doing a field test for T1 and T2, dynamic heart rate zones can be set as shown in Figure 5. Threshold Training System.

HEART ZONE		ZONES	CHAR	Т	CÝ	CHUSON
Zone Number	Zone Name	Zone Description	Zone Calories Burned/minute	Zone Sustainable Workout Time	Training Load* Points/Minute	Zone Benefits
×5	Very Hard Zone	High, hot, hard effort "I can't talk comfortably." Feels near max	> 15 calories	15 seconds - 40 minutes	6-8-10 points**	Get Fastest
		T2 HIGH T	HRESHOLD			
24	High Moderate Zone	Hard strong effort "I can talk but it is not comfortable." Feel shift in breathing rate	10-14 calories	40 minutes - 2 hours	4 points	Get Fast
Na S	Low Moderate Zone	Moderate effort "I can talk conversationally." Feels sustainable	7-9 calories	1-3 hours	3 points	Get Fitter
		T1 LOW T	HRESHOLD			
M2	High Easy Zone	Easy effort "I can talk and it is comfortable." Feels comfortable-enjoyable	4-6 calories	Very, very long time - possibly "all day"	2 points	Get Fit
ZONE	Low Easy Zone	Very easy effort "I can talk and it is easy." Feels almost effortless	1-3 calories	Long-lasting	1 points	Get Healthy

Figure 5. Threshold Training System Zones Chart Using Low and High Threshold Heart Rate Numbers to Anchor Zones (Edwards)

Conclusion

The MHR 220-minus-age equation – still in use in health clubs, schools, medical settings, and among many coaches today – has been proven to be outdated and inaccurate. Mark Sarzynski, PhD, a prolific researcher in personalized exercise medicine at the Arnold School of Public Health at the University of South Carolina's concluded, "Our results fail to validate the effectiveness of either of the

¹⁹ Giancarlo Condello,1, Ezekiel Reynolds,2, Carl Foster,2,3,Jos J. de Koning,2,3, Erika Casolino,1, Megan Knutson,2, and John P. Porcari2, A Simplified Approach for Estimating the Ventilatory and Respiratory Compensation Thresholds J Sports Sci Med. 2014 May; 13(2): 309–314. 2014 May 1.

two most widely used age-based HRmax prediction equations in sedentary, healthy adults. These results suggest that it may be very difficult, perhaps even impossible, to predict with a low standard estimate of error HRmax from age.²⁰

That certainly has been the conclusion – and experience – of Carl Foster, the former collegiate cross country runner with the mere 160 BPM MHR who now is Professor of Exercise Science at the University of Wisconsin, La Crosse as well as the past editor of the International Journal of Sports Physiology and Performance. "There is not one equation that is meaningful or any better than any other equation," he wrote, "except for some very specific uses in medical practice, I think that the use of any predicted MHR should not be used in the exercise industry, in schools, and for other uses because it is useless at best, and dangerous at worst."

Using equations that predict MHR individually is a fruitless process because individuals simply have too much variance in MHR. Rather, an individual's MHR should be estimated using sub-max field tests with results used to derive training zones specific to the individual as well as help in the development of safe and appropriate exercise prescriptions for individuals.

There is no reason to continue using the 45-year old MHR 220-minus-age equation given what we know today.

She can be reached at (1) 916.481.7283 or by email <u>sally.edwards@heartzones.com</u>. She lives and works in Sacramento, California.

Sally Edwards, MA, MBA, Physical Educator, is the CEO and founder of Heart Zones, Inc, the fitness technology company for Smart health and fitness. She is the author of 25 books on exercise, triathlon and running, and weight loss. She currently develops heart rate technology and programs for schools, health clubs, corporate wellness, sports teams, and rehabilitation centers.

Ms.Edwards has been an avid heart rate monitor user since she qualified and competed in the 1984 Olympic Trials guided by her Polar© heart rate monitor 32-years ago. Her oft-stated goal in life is to get America fit and one way is to use wearables like a heart rate monitor. She developed the Blink line of heart rate monitors and sensors for her program, Heart Zones Training[™]. A professional runner and triathlete, she first established the first five zone MHR system in 1992 in her first book on using a wearable for health and fitness *The Heart Rate Monitor Book*. She continues today to debunk the myth that MHR can be determined by any mathematical equation.

²⁰ Ibid M.A. SARZYNSKI

ADDENDUM 1

for PHYSICAL EDUCATION AND HEALTH TEACHERS

White Paper: The One-Size-Fits-All Age Adjusted Maximum Heart Rate Equation Fits No One

Change Physical Education Programs for Our Kids - The Smart PE Movement

At the request of physical education and health teachers in schools in which Heart Zones, Inc. has installed the company's group heart rate system, The Heart Zones System²¹, I have been asked to make several suggestions about the use of the Maximum Heart Rate (MHR) equation 220 minus age for children and adolescents in Physical Education classes.

Often, PE teachers grade students based on their time in a "target zone," which this White Paper demonstrates is unfair because it uses the one-size-fits-all calculations. Each student's MHR is different and as a result every student's training zones are different. Just as in academic courses like reading, in which each student reading level and skills are individually assessed,²² physical education and health classes should follow these best practices. Each student should take a field test for either Threshold heart rates or MHR and use personalized zones that are safe and appropriate for them. It simply isn't fair to a low MHR student, which in this White Paper is like a blue whale kid, or a high MHR student or referenced as a hummingbird kid with a fast MHR to have a fixed target heart rate zone based on their age. Not all 10-year old students have a MHR of 210 BPM or 220 minus the age of ten.

We recommend that PE and health teachers follow best practices by using objective and individualized grading. The typical field test for assessing MHR²³ takes no more than 20 minutes and can easily be conducted within a single PE class period. For more information and a complimentary lesson plan for assessing MHR or Threshold zones, please contact Heart Zones, Inc. by email at staff@heartzones.com

²¹ http://schools.heartzones.com/

²² Arq Bras Cardiol. 2011 Aug;97(2):136-40. Epub 2011 Jul 1.Validity of MHR prediction equations for children and adolescents. Machado FA1, Denadai BS.

²³ The Heart Zones Modified Pacer Test© is a field test for assessing MHR number and threshold heart rate for students.